

Notes

Indirect complexometric determination of mercury(II) using 1-propane thiol (PT) as a selective masking agent

P Parameshwara^a, J Karthikeyan^a, A Nityananda Shetty^{a*} & Prakash Shetty^b

^aDepartment of Chemistry, National Institute of Technology
Karnataka, Surathkal
Srinivasnagar 575 025, India

^bDepartment of Chemistry, Manipal Institute of Technology
Manipal 576 119, India

Received 9 February 2005; revised received 10 August 2005;
accepted 8 September 2005

A complexometric method for the determination of mercury(II) in presence of other metal ions is described based on the selective masking action of 1-propane thiol towards mercury(II). Mercury(II) present in a given sample solution is first complexed with an excess of EDTA and the unreacted EDTA is titrated against lead nitrate solution at pH 5-6 (hexamine buffer) using xylenol orange as the indicator. A 0.8% aqueous solution of 1-propane thiol is then added to displace EDTA from the Hg(II)-EDTA complex. The released EDTA is titrated with standard lead nitrate solution as before. Reproducible and accurate results are obtained for 4-85 mg of mercury(II), with relative error less than ± 0.23 % and coefficient of variation not more than ± 0.41 %. The effects of various ions were studied. The method is used for the analysis of mercury(II) in its synthetic alloy mixtures and also in complexes.

Keywords: Mercury(II), 1-propane thiol, masking agent, complexometric titration

IPC Code: C22B43/00

Mercury plays an important role in chemical and biological processes. Because of the extensive applications and toxic nature of mercury compounds, there is a need for a simple and rapid analytical method for the determination of mercury in samples.

Mercury(II) is not determined normally by direct EDTA titration, particularly in the presence of other metal ions¹ as EDTA is an unselective complexing agent, forms stable complexes with most of the metal ions. Usual practice is to complex mercury(II) together with the associated metal ions by EDTA and then dissociate selectively the Hg(II)-EDTA complex with an appropriate masking agent. The released

EDTA is titrated with standard metal ion solution. Singh² has described the determination of mercury(II) in the presence of various cations with thiourea as masking agent. In this method, the interference of copper(II) is avoided by controlling the pH at 5.5 followed by cooling the solution at 15°C before the addition of thiourea. Good results in the presence of copper(II) were obtained with thiourea as masking agent when it was present in slight excess over the required amount. This causes problem when a sample of unknown composition needs to be analysed. Selective determination of mercury using *N*-allylthiourea³ as masking agent requires heating to decompose the Hg-EDTA complex and some precipitation of HgS is also obtained. In the selective determination of mercury using thiosemicarbazide⁴, as masking agent, copper causes serious interference. Ueno⁵ suggested potassium iodide as masking agent in alkaline medium for determining mercury in the presence of copper, but many other cations interfered. 2-Mercaptoethanol⁶, 3-mercapto-1,2-propanediol⁷, 1,10-phenanthroline⁸, DL-cystein⁹, cysteamine hydrochloride¹⁰, thioglycolic acid¹¹, potassium bromide¹², glutathione¹³ were also used as a selective masking agent for the determination of mercury(II). Some of the other masking agents such as 4-amino-5-mercapto-3-n-propyl-1,2,4-triazole¹⁴, 2-imidazolidinethione¹⁵, hexahydropyrimidine-2-thione¹⁶ require tedious and time consuming synthesis procedures for their preparation, as they are not readily available.

The present investigation describes the use of 1-propane thiol as a masking agent for the selective and quantitative determination of mercury(II). The effects of foreign ions are studied and the applications of the method in the analysis of mixture of ions and mercury complexes are also reported.

Experimental Procedure

All reagents used were of analytical or chemically pure grade. The stock solutions of mercury(II) chloride, EDTA (~ 0.04 M) (sodium salt), and lead nitrate (0.02 M) were prepared by dissolving a requisite amount of salts in minimum amount of water, making up to the mark with distilled water and standardizing the solution by the standard methods¹⁷. Freshly pre-

*For correspondence (E-mail: nityashreya@rediffmail.com; Fax: +91-824-2476090)

pared (0.5 %) aqueous solutions of xylenol orange (indicator) and (0.8 %) 1-propane thiol (masking agent) were prepared by dissolving the requisite amount of reagent in distilled water. Solutions of various metal ions were prepared by dissolving calculated amounts of the metal chlorides/nitrates/sulphates in distilled water or with suitable acids and then making up to a known volume.

Method

To an aliquot of sample solution containing 4-85 mg of mercury(II) and varying amounts of diverse metal ions, an excess of 0.04 M EDTA was added and the solution was diluted with 25mL of distilled water. The pH of the solution was adjusted to 5-6 by adding solid hexamine. The surplus EDTA was back titrated with standard lead nitrate solution to a sharp colour change of xylenol orange from yellow to red. To this, a freshly prepared 0.8% solution of 1-propane thiol(PT) was added in required amount. The contents were mixed well and allowed to stand for 5 min in order to ensure the quantitative release of EDTA. The liberated EDTA was then titrated with the standard lead nitrate solution as before. The second titre value is equivalent to the amount of mercury(II) present in the aliquot.

Analysis of mercury complexes

Mercury complexes with thiourea, thiocyanate, 1,2,4-triazole-3(5)-thiol, thiocarbohydrazide were prepared and purified by the reported methods¹⁸⁻²². A known weight (approx. 1g) of the complex was carefully decomposed with aqua regia by evaporation to near dryness. The residue was then cooled, dissolved in distilled water and made up to a 100 mL standard flask. Aliquots of this solution were used for the estimation of Hg(II) as per the proposed method.

Results and Discussion

Masking action of 1-propane thiol

1-Propane thiol(PT) acts as a monodentate ligand and forms a stable complex with mercury(II). According to HSAB theory^{23,24}, mercury(II) forms strong bond through soft sulphur of mercapto group. Therefore, it is reasonable to expect the bonding of Hg(II) with deprotonated sulphur of thiol group. The quantitative release of EDTA from Hg-EDTA complex by 1-propane thiol indicates that Hg-PT is more stable than Hg-EDTA complex under the conditions employed. The release of EDTA is quantitative and instantaneous at 25-30°C itself. The Hg-PT complex

formed is soluble under the experimental conditions and the detection of the end point is very sharp.

Effect of PT concentration

It was observed that for instantaneous and quantitative release of EDTA from the Hg(II)-EDTA complex, the amount of PT required was in the molar ratio of 1:5 (M:L). Further, it was noticed that the addition of excess PT, as much as 20-fold excess over the required molar ratio does not have adverse effect on the results obtained. In all subsequent determinations, the concentration of PT was maintained at slight excess over the 1:5 (M : L) molar ratio.

Accuracy and precision

In order to check the accuracy and precision of the method, determination of mercury in the concentration range 4-85 mg were carried out under the optimized experimental conditions. These results are presented in Table 1. The results show that the maximum relative error and coefficient of variation (n=6) of the method are $\pm 0.23\%$ and $\pm 0.41\%$, respectively. From these results, it is reasonable to infer that the proposed method is precise and accurate.

Effect of foreign ions

The effect of various cations and anions on the quantitative determination of Hg(II) was studied by estimating 21.40 mg of Hg(II) in the presence of different metal ions. No interference was observed for the following ions at the amounts in mg shown: K(50), Zn(II) (250), Pb(II) (220), Cd(II), La(II) (100), Ba(II) (60), Co(II) , Ni(II) (50), Cu(II) (30), Mn(II) (10), La(III)(100),Y(III) (50), Ir(III), Rh(III) (30), Al(III), Au(III) (20), Bi(III) (10), Ce(III), Ru(III) (5) , Pt(IV) (30), As(IV) (25), Se(IV) (5), U(VI) (25), acetate(140), borate(200), chloride(150), sulphate, tartar-

Table 1—Precision and accuracy in the determination of mercury(II)

Mercury, mg		Relative error (%)	Standard deviation (mg)	Coefficient of variation (%)
Taken	Found			
4.28	4.29	+0.23	0.04	0.33
8.56	8.57	+0.12	0.04	0.41
12.84	12.83	-0.08	0.05	0.36
17.12	17.10	-0.12	0.05	0.26
21.40	21.40	0.00	0.01	0.05
25.68	25.68	0.00	0.05	0.18
34.24	34.22	-0.06	0.05	0.13
42.80	42.79	-0.02	0.01	0.03
64.20	64.19	-0.02	0.01	0.03
85.60	85.55	-0.06	0.02	0.03

Table 2— Analysis of mercury complexes (n=3)

Complex	Hg (II) calculated (%)	Hg (II) found (%)	Relative error (%)
Hg(CH ₄ N ₂ S)Cl ₂ .1/2 H ₂ O ^a	56.69	56.61	-0.14
Hg(CH ₄ N ₂ S) ₂ Cl ₂ ^b	47.34	47.40	+0.13
Hg(CH ₆ N ₄ S) ₂ Cl ₂ ^c	41.46	41.40	-0.15
Hg[Zn(SCN) ₄] ^d	40.27	40.20	-0.17
Hg(C ₂ H ₂ N ₃ S) ₂ ^d	50.05	49.96	-0.18

^aMercury complex of thiourea^bMercury complex of thiourea^cMercury complex of thiocarbohydrazide^dMercury complex of 1,2, 4-triazole-3(5)-thiol

Table 3— Determination of mercury(II) in synthetic mixtures(n=5)

Mixture	Composition (mg)	Hg (II) found (mg)
Hg + Zn + Pb	21.4 + 50.0 + 80.0	21.38
Hg + Co + Cd	21.4 + 40.0 + 40.0	21.43
Hg + Zn + Ni	21.4 + 50.0 + 30.0	21.42
Hg + Zn + Pb + Ni	21.4 + 50.0 + 60.0 + 30.0	21.43
Hg + Zn + Ni + Cd	21.4 + 60.0 + 40.0 + 30.0	21.38
Hg + Zn + Ni + Cd	14.2 + 39.6 + 26.4 + 19.8	21.42

rate (250), oxalate(50), phosphate(100). Metal ions such as Pd(II) as its chloride, Tl(III) as its nitrate, Cr(III) and Sn(IV) as their chlorides show severe interference giving positive errors. This is due to the release of EDTA from their respective complexes by the reagent. However, the interference of Pd(II) (up to 5 mg) and Tl(III) (20 mg) can be avoided by premasking these ions with L-histidine (5% C₃H₉N₃O₂, 15-20mL) (0.75 - 1.0 g) and hydrazine sulphate (5% N₂H₄.H₂SO₄, 10mL) (0.5 g), respectively. The interference of Cr(III) is due to the deep purple colour of its EDTA complex, which makes the detection of the end point rather difficult.

Applications

In order to explore the practical application of the proposed method, it was extended for the determination of mercury in its complexes and in synthetic mixture of metal ions. The experimental results of these analyses are presented in Tables 2 and 3, respectively.

It is evident from these results that the method can be conveniently employed in the analysis of mercury in its complexes and alloys with a fair degree of accuracy.

Conclusions

The proposed method is simple as it does not require any adjustment of pH after the addition of the reagent or heating for the release of EDTA from the Hg(II)-EDTA complex. Since many metal ions and anions do not show interference, the method is fairly selective for the rapid analysis of mercury in the presence of these ions.

Reference

- West T S, *Complexometry with EDTA and Related Reagents* (BDH, Poole), 1969, 105.
- Singh R P, *Talanta*, 16 (1969) 1447.
- Vasilikiotis G S & Apostolopoulou C D, *Micro Chem J*, 20 (1975) 66.
- Korbl J & Pribil R, *Chem Listy*, 51 (1957) 667.
- Ueno K, *Anal Chem*, 29 (1957) 1669.
- Muralidhara Rao B & Narayana B, *Chim Acta Turc*, 21 (1993) 27.
- Prakash Shetty, Khader A M A, Nityananda Shetty A & Gadag R V, *Rev Roum Chimie*, 40(4) (1995) 351.
- Ramachandra B & Narayana B, *Ann Chim*, 87 (1997) 647.
- Abraham Joseph & Narayana B, *Res J Chem Environ*, 3 (1999) 49.
- Abraham Joseph, Narayana B & Bhat K S, *J Indian Chem Soc*, 76 (1999) 510.
- Prakash Shetty, Nityananda Shetty A & Gadag R V, *Rev Roum Chimie*, 45(9) (2000) 841.
- Sree Kumar N V, Nazareth R A, Narayana B, Prashantha Hegde & Manjunatha, *Micro Chim Acta*, 140 (2002) 63.
- Prakash Shetty & Nityananda Shetty A, *Turk J Chem*, 28 (2004) 573.
- Gadiyar H R A, Gadag R V & Gajendragad M R, *Talanta*, 29 (1982) 941.
- Narayana B & Gajendragad M R, *Talanta*, 35 (1988) 719.
- Narayana B, Kuchinad G T & Gajendragad M R, *Chinese J Chem*, 10 (1992) 394.
- Vogel A I, *A Text Book of Quantitative Inorganic Analysis*, 3rd edn (Longmann, London), 1964.
- Rosenheim A & Meyer V J, *Z Anorg Chem*, 49 (1906) 13.
- Malay R, *Ber*, 9 (1876) 172.
- Claus A, *Ber*, 9 (1876) 226.
- Burns G R, *Inorg Chem*, 7 (1968) 277.
- Gupta B K, Gupta D S, Dikshit S K & Agarwala U, *Indian J Chem*, 15A (1977) 624.
- Perarson R G, *Chem Eng News*, 43 (1965) 90.
- Perarson R G, *Chem Brit*, 3 (1967) 103.